

Pachauri, Saroj.: Unmet Reproductive and Sexual Health Needs in South Asia. The Journal of Family Welfare. Journal of Health & Population in Developing Countries. 1998.1(2).p.29-39.

Unmet Reproductive and Sexual Health Needs in South Asia

Saroj Pachauri

The magnitude of reproductive and sexual health problems in South Asia is daunting. However, an enabling policy environment provides an opportunity to address unmet needs. Neglected reproductive health problems can be effectively addressed through a life-cycle approach. The paper underscores inter-linked gender, sexuality and rights concerns within which these problems are embedded.

Global Consensus on Reproductive and Sexual Health

At the 1994 International Conference on Population and Development (ICPD) in Cairo, the world community effectively wrote a new agenda for population and development to serve as the blueprint for population programs internationally and nationally for the next twenty years. ICPD placed the population problem squarely within the development context focusing particularly on quality of life, human rights and women's empowerment.

At Cairo, the nations of the world agreed that governments should give special attention to the health of women, the survival of infants and young children, the education of girls, and in general, the empowerment of women. At the same time they should provide comprehensive reproductive health services to enable couples to achieve their reproductive goals, and determine freely and responsibly the number and spacing of their children. The ICPD consensus implies that if governments ensure that this basic package of social policies and reproductive health services is in place they will simultaneously make strides toward greater social equity and reducing high rates of population growth Sinding and Fathalla.

Consensus on the ICPD Program of Action provides the foundation for practical progress towards a number of interrelated sustainable development objectives.

However, to achieve these objectives, many unmet needs must be addressed. Several of these can be effectively met only if:

- Population agencies move beyond family planning as their main program mechanism and collaborate much more closely with health and development agencies;
- Development sectors integrate population related and human rights concerns into their policies and programs;
- Communities and governments come to terms with imbalances in power and opportunities between the state and its people, between providers and users of services and between men and women; and
- Governments foster participatory processes that give leadership and responsibility to communities, non-government organizations (NGOs), and the larger civil society.

Translating Reproductive and Sexual Health Rhetoric into Reality

While the global acknowledgment of reproductive and sexual health has brought much satisfaction and sense of achievement to many who worked for well over a decade to help develop the new paradigm articulated at ICPD, it has also thrown up fresh challenges. A major challenge confronting governments now is how to translate reproductive and sexual health concepts into policies and programs within national contexts.

To translate reproductive rhetoric into reality, two-priority issues must be addressed. First, it is essential to make a conceptual shift. Second, packages of good quality services must be designed and implemented to address reproductive and sexual health needs of clients. Clients have multiple needs, many of which are unmet, especially in South Asia, a particularly deprived region. An important priority is to design programs that are truly responsive to the needs of the clients. In this paper unmet reproductive and sexual health needs in South Asia are defined and discussed with some reference to programs. A discussion of implementation of program strategies to address these needs is not within the scope of this analysis.

South Asia within a Global Context

With a share of 22 percent of the world's people, South Asia is inhabited by 40 percent of the world's absolute poor and is now the poorest region of the world. Its per capita income of \$309 is far below that of \$555 of Sub-Saharan Africa and \$970 for all developing countries. Its adult literacy of 48 percent is below 55 percent of Sub-Saharan Africa and 98 percent of East Asia (excluding China) (Table 1). About one-half of the world's illiterate people live in South Asia. Neglect of basic social services is related to poverty in South Asia. The World Bank estimates that some this region contains approximately 40 percent of the world's absolute poor surviving on less than one dollar a day. Far more crippling than income poverty is the poverty of basic human capabilities, which prevents people from taking advantage of market opportunities. Nearly two-thirds of the population in South Asia is deprived of basic human capabilities, compared to just over one-quarter, which is deprived of a minimum income. This makes the task of poverty elimination doubly difficult UNDP.

The impact of deprivation is greatest on women and children. The maternal mortality rate in South Asia is about 600 per 100,000 live births. It is as high as 850 in Bangladesh, 1,500 in Nepal, and 1,600 in Bhutan. These are among the highest maternal mortality rates in the world, when compared to a rate of only 28 in industrial countries. Despite its much higher GNP growth rate and its more robust increase in food production. South Asia has more malnourished children than Sub-Saharan Africa. Almost, two-thirds of children in South-Asia are underweight, compared to one-sixth in Sub-Saharan Africa. Approximately one-third of all babies in India and one-half in Bangladesh are born with low birth weight. In Sub-Saharan Africa, the proportion is about one-sixth UNICEF.

Table 1: Comparative Education and Health Profile for South Asia, East Asia, Sub-Saharan Africa and Developing Countries.

	South Asia	East Asia (excl. China)	Sub-Saharan Africa	All Developing Countries
Education Profile				
Adult literacy (%) 1993 1 st , 2 nd and 3 rd level gross	34	94	45	60
Enrollment ratio (%) 1993	43	76	37	51

Mean years of schooling 1992	1.2	6.2	1.3	3.0
Health Profile				
Female life expectancy 1993 Maternal mortality rate per	61	75	53	63
100,000 live births 1993	585	100	929	384
Total fertility rate 1993	4.2	1.9	6.4	3.6
Women using contraception (%) 1986-93	39	79	15	55

Source : Haq ul 1997.

Defining Unmet Need

Traditionally, the concept of unmet need originates from the field of family planning where it has been extensively used to measure the proportion of women/couples who want to limit and/or space their births but are not using any form of contraception. Because family planning is now seen as an integral part of reproductive and sexual health, this concept of unmet need must be redefined. In a recent essay, Ruth Dixon-Mueller and Adrienne Germain define a broader scope of unmet need. They argue that the concept of unmet need should include:

- Recognizing the need among non-users at risk of unwanted pregnancy for any method of contraception;
- The need among some users for a more effective, satisfactory, or safer method;
- The need among both users and non-users for treatment of contraceptive failure (or non-use) through safe and accessible abortion services; and
- The need for related reproductive health services such as the prevention and treatment of reproductive tract infections (RTIs) Germain and Dixon-Mueller.

The concept of unmet need is likely to evolve and be further refined as programs make a paradigm shift and begin to implement comprehensive sexual and reproductive health services.

The Paradigm Shift

In order to operationalize reproductive health programs, a change in focus from a population, control approach of reducing numbers to a reproductive health approach of addressing the needs of clients is necessary. Implementing reproductive health services within national programs in South Asian countries may require an ideological shift. In India, for example, it would necessitate a change in the culture of the program which has focused in the past on achieving targets to one that will now aim at providing a range of good quality services. The implication is that reproductive health programs should be responsible for reducing the burden of unplanned and unwanted child bearing and related morbidity and mortality Jain and Bruce. Achieving the demographic goal of reducing the rate of population growth requires broader social and economic macro-level policies such as policies to improve education and enhance employment opportunities for women.

Reproductive Rights and Sexual Health

While reproductive health needs are beginning to be defined, those related to reproductive rights have barely begun to be even articulated in South Asia and are, therefore, largely unmet. The proponents of the reproductive health framework believe that reproductive health is inextricably linked to the subject of reproductive rights and freedom, and to women's status and empowerment. Thus, the reproductive health approach extends beyond the narrow confines of family planning to encompass all aspects of human sexuality and reproductive health needs during the various stages of the life cycle. In addressing the needs of women and men, such an approach places an emphasis on developing programs that enable clients to:

- Make informed reproductive choices and exercise their right to choose.

- Receive education and counseling services for responsible and healthy sexuality;

- Obtain services for the prevention of unwanted pregnancy, safe abortion, maternity care and child survival, and for the prevention and management of reproductive morbidity including gynecological problems, reproductive tract infections (RTIs), sexually transmitted infections (STIs) and HIV/AIDS.

Thus, reproductive and sexual health programs are concerned with a set of specific health problems, identifiable clusters of client groups, and distinctive goals and strategies.

Unmet Needs in South Asia

This evolving paradigm is being put to its most critical test in South Asia, which is fast emerging as the poorest, the most illiterate, the most malnourished, the least gender sensitive - indeed, the most deprived region in the world Haq ul (Table 1 & Table 2). A recent assessment of human development in South Asia, including India, Pakistan, Bangladesh, Nepal, Sri Lanka Bhutan, and Maldives, showed Maldives, showed that while these seven countries share many common characteristics, their performance in raising the level of human development is variable. There has been some progress in the field of health in South Asia. Infant mortality has been cut in half in the last three decades and life expectancy has increased by 17 years, from 44 in 1960 to 61 in 1993. But, there are serious gaps in the field of education. More than, one-half of the population in South Asia is still illiterate.

Table 2: Selected Health Indicators for Countries in South Asia and Developing Countries.

	India	Pakistan	Bangladesh	Nepal	Sri Lanka	South Asia Weighted Average	Developing countries
Population with access to health services (%) 1993	85	55	45	N/A	93	78	80
Population per doctor	3,140	3,780	11,350	38,650	6,230	4,665	N/A
1976	2,439	2,000	12,500	16,667	7,143	3,697	5,767
1988-91							

Population per nurse	6,320	10,040	53,700	52,770	2,240	12,061	N/A
1976	3,333	3,448	20,000	33,333	1,745	5,465	4,715
1988-91							
Maternal mortality rate per 100,000 live births 1993	570	340	850	1,500	140	583	384

Source: Haq ul. 1997.

Human deprivation is especially grim for women. Two thirds of adult women in the region are illiterate and maternal mortality is among the highest in the world. A comparison of South Asia's performance with that of East Asia is revealing. Starting with similar income levels three decades ago, East Asia is now way ahead of South Asia in its social and health indicators. It is clear that South Asia has failed to provide adequate coverage of basic social services to its 1.2 billion people (Haq ul).

Although a sub-continent, there are significant differentials among the countries of South Asia (Table 3 & Table 4). At one end of the spectrum is Sri Lanka - its human development and health indicators are among the highest in the world, often surpassing those of many developed countries. Its adult literacy, for example, is 90 percent. Population growth rate is 1.5 percent, compared to an average of 2.3 percent for South Asia. Its life expectancy at 72 is 11 years longer than the South Asian average of 61 years. Infant mortality in Sri Lanka is 15 per 1000 live births compared to 91 in Bangladesh, 95 in Pakistan, and 79 in India. Basic health facilities are available to 93 percent of Sri Lanka's population (Haq ul). In contrast, other South Asian countries, such as Pakistan, India, and Bangladesh depict gloomy socio-demographic and health profiles indicating a huge unmet need in these countries. The Reproductive Risk Index designed by Population Action International ranks India and Sri Lanka in the moderate risk category, and Bangladesh, Nepal, and Pakistan in the high risk group Population Action International.

Table 3: Selected Social Health Indicators for Countries in South Asia.

	India	Pakistan	Bangladesh	Nepal	Sri Lanka
Population in absolute	416	36	60	8.4	6.0

poverty (%) 1993 number (millions) as % of total population	46	28	52	40.0	31.0
Population without access to health services 1993 number (millions) as % of total population	135 15	60 45	63 55	N/A N/A	1.3 7.0
Illiterate female adults 1993 number (millions) as % of total adult female population	183 64	28 77	26 75	5.4 87.0	0.8 14.0
Malnourished children under 5, 1994 number (millions) as % of total under-5 population	62 53	10 40	11 67	2.0 49.0	0.8 38.0
Under-five mortality rate 1994 (per 1,000 live births)	119	137	117	118	19

Source: Haq ul. 1997.

Table 4: Profile of Human Development for Countries in South Asia.

Country	Life expectancy at birth (years) 1994	Population with access to			Daily calorie supply per capital 1992	Adult literacy rate (%) 1994
		Health services (%) 1990-95	Safe water (%) 1990-95	Sanitation (%) 1990-96		
Thailand	69.5	90	89	96	2,443	93.5
Philippines	67.0	71	86	77	2,258	94.4

Indonesia	63.5	93	62	51	2,755	83.2
Vietnam	66.0	90	43	22	2,250	93.0
Myanmar	58.4	60	60	43	2,598	82.7
India	61.3	85	81	29	2,395	51.2
Pakistan	62.3	55	74	47	2,316	37.1
Bangladesh	56.4	45	97	48	2,019	37.3

Source: Haq ul. 1997.

Addressing Unmet Needs

Addressing unmet needs implies designing strategies to target neglected client groups such as women, men and adolescents as well as implementing services to meet their priority reproductive and sexual health needs throughout the life cycle.

Unmet Needs of Women: Users' Perspectives on Improving Quality of Care

Past programs have focused primarily on addressing family planning needs. Future programs must also address other reproductive health needs of women. Family planning efforts should be focused on: (1) reaching the unreached especially where access remains a problem as in Nepal, Pakistan, and Myanmar and serving unserved areas of other countries; (2) expanding contraceptive choice, particularly in countries like India, where the program essentially relies on a single method; and (3;) improving the quality of services, which is an unmet need of all countries in the region.

If reproductive health programs are designed to address client's needs, the quality of services must be improved, particularly is perceived by clients. There is a need to focus on women since they constitute the major client group or users of these programs and also have the greatest problem of access, both physical and social, to health services Pachauri .

There are ample data worldwide showing a high burden of reproductive morbidity among women in developing countries. According to the World Bank, about one-third of the total disease burden in developing countries among women in the age group of 15 to 44 years is linked to health problems related to pregnancy, childbirth, abortion, human immuno-deficiency virus (HIV), and RTIs World Bank. Available data for South Asia show that women have a huge unmet need for services related to these conditions. Socio-economic and biological determinants operate synergistically throughout the lives of poor women in South Asia to undermine their health, resulting in high levels of morbidity and mortality Gittlesohn et al.; Akhtar, Rahman and Ahmed. However, among diseases for which cost-effective interventions exist, reproductive health problems account for the majority of the disease burden in women of this age group. There is, therefore, an urgent need to design and implement cost-effective programs to address women's reproductive health needs. Women-centered, gender-sensitive services must be organized and implemented (Pachauri 1995).

Since gender inequalities favor men in South Asian countries and sexual and reproductive health decisions are largely made by men, there is a growing realization that unless men are reached, program efforts will have limited impact. Research on sexuality, especially in the field of HIV/AIDS, has highlighted the inadequacy of strategies that target only women. Because of the prevailing gender power equation, women are not only especially vulnerable, but are also unable to negotiate changes in sexual behavior. Research on sexual negotiation has dramatically underscored the need for involving men in programs that aim at bringing about changes in sexual behavior for the prevention of infection. However, such behavioral change is relevant not only for the prevention of infection but also for addressing other reproductive and sexual health problems. Reproductive and sexual health needs of men have not been addressed by past programs that have mainly targeted women. Men too have unmet needs as illustrated by, recent work in India Khan, Khan and Mukherjee and Bangladesh Hawk. Therefore, the involvement, of men as partners is essential Pachauri.

Unmet, Needs of Adolescents: A Neglected Group

Adolescents are another neglected client group. Adolescents have been by passed by all programs and consequently their needs have neither been assessed nor addressed. Health services for the adolescent girl have special significance in South Asian countries where there is a strong son preference. Such services would not only affect the health of the adolescent girl but would also have long-term inter-generational effects by reducing the risk of low birth weight and

minimizing subsequent child mortality risks. So far, neither services nor research have focused on the adolescent's health and information needs. Youth 10-24 years of age represent about a third of the population in the countries of South Asia Population Reference Bureau. Therefore, the consequences of this neglect takes on enormous proportions, particularly as adolescents are exposed to the rising threat of HIV and STIs (Pachauri).

Reducing the Burden of Reproductive and Sexual Health Morbidity

Data on the magnitude of reproductive and sexual health morbidity in South Asia are sparse. Research is urgently needed to systematically assess the extent of reproductive and sexual health problems in the countries of the region so that program priorities can be delineated. The following section provides an assessment of the magnitude of priority problems, which must be urgently addressed.

Maternal Mortality and Morbidity: A Neglected Tragedy

The magnitude of women's reproductive health problems is reflected in the number of deaths related to pregnancy and childbirth, the most direct indicator of reproductive health care. Worldwide, there are half a million deaths of women each year related to pregnancy, and all but 6,000 of these occur in the developing world. Maternal mortality is notoriously difficult to measure accurately and so only estimates are available. India's maternal mortality ratio, 400/500 per 100,000 live births, is fifty times higher than that of many developed countries and six times higher than that of Sri Lanka Ascadi and Johnson-Ascadi. The maternal mortality ratio is estimated at 450 in Bangladesh (Ascadi and Johnson-Ascadi 1990) and at 1,500 in Nepal per 100,000 live births United Nations.

Mortality statistics, however, tell us only, a part of the story. For every woman who dies, many more suffer serious illness. Community based data on maternal morbidity are scarce. In recent studies, for every maternal death there were 643 morbidity's in Bangladesh and 541 in India Fortney and Jason. In Bangladesh and India, 80 and 58 percent women respectively, reported at least one morbidity, during pregnancy and the puerperium; 24 and 2 percent respectively, reported five or more morbidities. A life threatening morbidity was reported by 5 percent of Indian women and 32 percent of Bangladeshi women. Life threatening and serious morbidities were reported by 46 percent of Indian and 65 percent Bangladeshi women. For every woman who died from pregnancy-related causes,

65 percent women in Bangladesh and 25 percent in India suffered from chronic morbidity (Fortney and Jason (eds) 1996).

These data on maternal mortality and morbidity underscore the high level of unmet need for maternity care among women in South Asian countries. By some estimates, better care during labor and delivery could prevent 50-80 percent of maternal deaths Rodriguez et al.; Walker et al.. But, the vast majority of births in India, Bangladesh, and Pakistan take place at home and, are handled mostly by untrained-birth attendants. As recently: as 1992-1993, no more than 16 percent of all rural births in India were conducted in institutions and as many as two-thirds were delivered by traditional birth attendants International Institute of Population Studies. In Sri Lanka, however, trained personnel attend 94 percent of the births United Nations.

Malnutrition is an underlying problem that seriously affects the health of adolescent girls and adult women and has its roots in early childhood. About 85 percent women in South Asia are anemic. The association between anemia and low birth weight, prematurity, perinatal mortality and maternal mortality has been extensively documented in South Asian countries.

An area that has not been explored relates to violence and injuries, which accounted for 14 percent of maternal deaths in a study in Bangladesh Rahman, Whittaker and Hossain. Recent research shows that gender violence is, pervasive in South Asian countries Sidney, Hashemi and Riley; Sood (ed); Deraniyagala. This problem, however, remains invisible because of the silence that shrouds it. It is a problem that is denied by society.

Unsafe Abortion: A Pervasive Problem

Unsafe induced abortion is the greatest single cause of mortality, and at the same time, the most preventable one. Of all the major causes of maternal deaths, those that lead to abortion deaths are the best understood. Women do not need to die or suffer medical consequences from abortions because abortions do not necessarily kill; it is carelessly performed abortions, which kill Maine. Abortion is illegal in most South Asian countries. In India abortion was legalized over twenty years ago, but is still a neglected problem. Access to safe abortion services for poor women, especially in rural areas, remains problematic in India.

The conceptual link between family planning and abortion is fundamental. Effective contraception is an important means of preventing unwanted pregnancy thus pre-empting the need for abortion. In the absence of safe contraceptive backup, however, women will continue to be forced to employ unsafe means for terminating unwanted pregnancies with attendant high maternal mortality and morbidity.

Data on abortion are unreliable and difficult to obtain. According to official estimates 1.5 million abortions occur in South Asia each year. The estimated abortion rates per 1000 women of reproductive age are 3.4 in Bangladesh, 3.0 in India, 4.1 in Nepal, and 5.4 in Sri Lanka International Planned Parenthood Federation South Asia Region. These figures are likely to be significant underestimates because abortion is illegal and/or severely restricted in all South Asian countries except India. Therefore, the majority of abortions in the region are clandestine; are usually performed under unsafe conditions; and are not reported Kapoor.

Reproductive Tract Infections: A Silent Epidemic

Since community-based research on women's reproductive illnesses, including RTIs and STIs, was neglected in the past, these problems remained invisible Bang and Bang. Rani Bang's landmark study in a rural, tribal area of Maharashtra, India, indicating that 92 percent of the women suffered from one or more gynecological problems and that the majority had never sought any treatment, created quite a stir nationally and internationally. This study not only depicted the magnitude of the problem of gynecological morbidity in poor women, but also brought into focus the neglect associated with these problems Bang et al. It highlighted the need for further research to assess levels of morbidity among poor women.

A recent review of nine community-based studies of gynecological morbidity in India showed a considerable burden of RTIs/STIs Koenig et al. In Maharashtra, the prevalence of infections including vaginitis, cervicitis, and pelvic inflammatory disease (PID) was 46 percent (Bang et al.). In four community-based studies of gynecological morbidity recently conducted in four different sites in India (rural West Bengal and Gujarat, and urban Baroda and Bombay), the prevalence of clinically diagnosed RTIs ranged from 19 to 71 percent Baroda Citizens Council. In a similar study conducted in rural Karnataka, over 70 percent of women had clinical or laboratory evidence of RTIs Bhatia et al. These studies leave little doubt that prevalence of RTIs in India is unacceptably high

and that gynecological morbidity constitutes a major public health problem among poor women.

Research in India shows that poor women carry a heavy burden of reproductive morbidity; a significant component of such morbidity is unrelated to pregnancy and is due to RTIs, many of which are sexually transmitted; these reproductive problems are invisible because of socio-cultural reasons and are unattended because women do not have access to health care for these illnesses Pachauai.

Studies in Bangladesh also show a significant burden of reproductive morbidity. In a survey of a rural community, where the magnitude and nature of morbidity due to RTIs among users and non-users of contraceptives was assessed, 22 percent of the 2,929 women surveyed reported symptoms of infection. Of the 472 symptomatic women examined, 68 percent had clinical or laboratory evidence of infection Wasserheit et al..

Thus, there is growing evidence that the burden of reproductive morbidity is unacceptably, high among poor women in South Asia. This represents an important unmet need because these RTIs are frequently asymptomatic in women and are not recognized and treated. Certain contraceptive methods increase the risks of RTIs. For example, oral contraceptives predispose women to candidiasis and chlamydia infection Wasserheit et al. These linkages between RTIs and contraception further complicate the problem, adding yet another dimension to the already complex challenge of addressing these problems.

Poor women are also at increased risk of RTIs because of the unhygienic management of menstruation. Childbirth and abortion exacerbate infections of the lower reproductive tract and facilitate their spread to the up-per tract. Untreated lower tract infections are likely to progress to pelvic inflammatory disease, which has serious sequelae including infertility, ectopic pregnancy; chronic pain and recurrent infection. The problem of infertility is particularly serious and can, be devastating for women in South Asia where child bearing is highly valued. RTIs and STIs, poor obstetric and gynecological practices, illegal abortion and postpartum and postabortion infections are all preventable causes of infertility (Pachauri). In addition, RTIs effect fetal wastage, low birth weight and congenital infection, and impact on pregnancy outcome, child health and survival. The treatment of these infections is, therefore, a cost-effective approach to reduce perinatal and neonatal mortality. Child survival programs currently underway in South Asia not designed to address these problems (Pachauri).

Cancers of the Reproductive System: An Un-addressed Problem So far, cancers of the reproductive tract have received little attention even though these increasingly contribute to reproductive morbidity in Cancer is one of the three leading causes of adult female mortality in both developed and developing countries. While breast cancer is the most frequently occurring cancer among women in the industrialized countries, cervical cancer is the most common malignancy in developing countries where it accounts for 20 to 50 percent of all cancers and 80 to 85 percent of all malignancies of the female genital tract Belsey and Royston. In India, the incidence of cervical cancer ranges from 15.4 to 46.5 per 100,000 women Indian Council of Medical Research.

There is an established association between RTIs and cervical cancer Mishra and Sinha; Indian Council of Medical Research N; Murthy, Sehgal and Satyanarayana. Early onset of sexual activity and multiple sexual partners increase the risk of cervical cancer Menon et al. The high incidence of cervicitis that is not due to conventional STI pathogens, has been postulated as a possible risk factor for cervical cancer among Indian women Luthra et al.

Cervical screening is an important means for preventing cancer but at present only limited screening facilities are available to women in South Asia. For example, while 15 percent of the world's cervical cancer cases exist in India, screening facilities are available only to a small minority of urban women.

HIV/AIDS: A Problem Denied

By the turn of the century, Asia is expected to become the epicenter of the global HIV/AIDS pandemic. Denial of the problem in South Asia continues to impede action. This represents a huge unmet need for South Asia.

After the first AIDS case was reported in India in May 1986, some 6,154 cases have been reported National AIDS Control Organization. These reported figures are believed to be a gross underestimate. Over the Years, there has been a steep rise in the reported seropositivity rate - from 2.5 per 1000 in 1986 to 20 per 1000 in 1998 National AIDS Control Organization. The prevalence of HIV infection has been on the rise in practically all states and all population groups in India. The increase is observed irrespective of when the infection reached a state. For example, no HIV infection was reported in Orissa till the end of 1992 but within a Year of its detection, prevalence increased about three-fold National AIDS Control Organization.

The northeastern region of India is experiencing a major HIV/AIDS epidemic. In this region, infection is spread predominantly by injecting drug users. The state of Manipur, which borders Myanmar, accounts for the highest number of cases in this region. After its initial detection in 1989, the prevalence of HIV among intravenous drug users in Manipur increased sharply. It was 64 percent by 1993 Sarkar et al. Manipur State borders the 'Golden Triangle' of Myanmar, Laos and Thailand and is on the drug-trading route. There is an explosive HIV/AIDS epidemic among drug users in Myanmar where over 60 percent of the world's opiates are produced.

The dominant mode of HIV transmission is heterosexual and this accounts for about two-thirds of the infections in South Asia. Therefore, addressing unmet need in the area of sexuality is a priority because changing sexual behaviors to promote safer sexual practices lies at the heart of HIV/AIDS prevention Pachauri. Information on HIV/AIDS and STIs in Pakistan, Nepal, and Bangladesh is inadequate and scanty. The magnitude of the problem is, therefore, not known. This implies that there is a serious unacknowledged and unmet need in these countries.

Integrating Services: A Bureaucratic Challenge

There is a growing realization that horizontal integration of services must be achieved if reproductive and sexual health and rights are to be universally realized. There must be a convergence of services at the user's level as the user or client has multiple reproductive health needs and often the same providers deliver services that are administered through multiple vertical programs. Vertical programs that originate from different government departments and are funded by multiple donors, each with its own agenda, can result in a multiplicity and fragmentation of services which can be wasteful and inefficient.

Thus far, most reproductive health programs are provided through vertical systems. Services for promoting family planning, improving child survival, and preventing HIV/AIDS are particular examples. It would be counterproductive to have reproductive health as yet another vertical program. In fact, the reproductive health approach provides an opportunity for integrating services. A critical mass of resources could also achieve greater cost-effectiveness.

in South Asian countries, governments have implemented family planning programs for the past three to four decades. Child survival programs have been implemented since the 1980s. Services for promoting safe motherhood have been put in place more recently. Prevention of HIV/AIDS and STIs are also more

recent initiatives. The challenge is to strengthen all these services by expanding their reach, improving their quality and effectively integrating additional reproductive health services within ongoing programs.

Gender Disparity: A Fundamental Problem

Women bear the greatest burden of human deprivation in South Asia. The Gender-related Development Index (GDI), which adjusts the measure of average human development to take account of gender disparities, shows a value of only 0.41 for South Asia, which is percent lower than the average for developing countries and less than half that of the industrial world. However, access to political, economic, and social opportunities, that is reflected by the Gender (GEM), starkly indicates deprivation among South Asian women. South Asia's GEM index of 0.23, is the lowest among all regions of the world, including Sub-Saharan Africa, (Haq ul).

For the most part, the neglect of women is related to low status and lack of autonomy. There is an overwhelming son preference in South Asian societies. South Asia is the only region in the world where men outnumber women. While Sri Lanka and Maldives have nearly equal male-female populations, the ratio of females for every 100 males is in Nepal, 93 in Pakistan, 94 in India (Haq ul). Some 74 million women are simply missing in South Asia! Women's voices have yet to be heard by the vanguards of society.

While the neglect of women is a widespread phenomenon throughout South Asia, there are some important differences among the countries of the region. Sri Lanka has performed fairly well in investing in and providing opportunities to women. In Sri Lanka, female literacy is 86 percent, female life expectancy is 106 percent of male life expectancy, and the female economic activity rate is 36 percent. Pakistan, on the other hand, ranks lowest in the region in most gender-related human development- indicators. Its female literacy rate is only 23 percent; female school enrolment is 16 percent; and female participation in economic activities is only 16 percent (Haq ul).

Concluding Comments

Even though reproductive-health rhetoric is now used by many, there are major information gaps at all levels ranging from a lack of understanding of the reproductive health and rights concepts to questions about what short-term and

long-term strategies are needed at the policy and program levels to implement services. This lack of information presents a major deterrent to: implementing programs to address unmet reproductive and sexual health needs in South Asia. Strong advocacy is needed to translate the rhetoric into reality. A range of different constituencies, including governments, donors, NGOs, activists, feminists, and researchers, must be informed and empowered to catalyze a process of growing number of stakeholders so that the ideology and the ethos embodied in the reproductive health concept effectively internalized and programs responsive to clients needs are designed and implemented (Pachauri 1995).

The magnitude of unmet reproductive and sexual health needs in South Asia is daunting. The problems in this deprived region are many. However, a changing policy environment presents an opportunity to make a difference.

Several countries in the region are making a paradigm shift and are redesigning their national programs to operationalize the ICPD Plan of Action. Reproductive and sexual health programs are now on the agenda of South Asia. India launched a major national Reproductive and Child Health program in October 1997. Bangladesh is in the process of designing a program Discussions are underway in other countries of the region to define reproductive health agendas and to redesign programs responsive to local needs.

Translating reproductive rhetoric into reality in South Asia presents a challenge that will require joint action by governments, donors, NGOs, the private sector and the larger civil society. Concerted efforts must be made to provide an enabling environment and to promote partnerships so that unmet reproductive and sexual health needs of the peoples of South Asia can be addressed by the involvement and participation of all.

References

[A] Akhtar, HH. MH Rehman and S Ahmed 1996. Reproductive Health, Issues and implementation Strategies in Bangladesh. Bangladesh Institute of Promotion of Essential and Reproductive Health and Technologies (BIRPERHT). Dhaka, Bangladesh.

[B] Ascadi. GTF and G Johnson-Ascadi 1990. Safe Motherhood in South Asia:

Socio-cultural and Demographic Aspects of Maternal Health. Background paper for the Safe Motherhood South Asia Conference, Lahore.

[C] Bang, RA and A Bang 1991 why Women Hide Them: Rural Women's Viewpoints on Reproductive Tract Infections. *Manushi* 69: 27-30.

[D] Bang. RA. AT Bang, M Baitule, Y Chaudhury, S Sarmukaddam and O Tale. 1989. High Prevalence of Gynecological Diseases in Rural Indian Women. *Lancet* 1(8629):85-88.

[E] Baroda Citizens Council Prevalence of Clinically Detectable Gynecological Morbidity in India. Results of Four Community-based Studies Unpublished manuscript.

[F] Belsey MA and E Royston. 1987, Overview of the Health of Women and Children. Technical background paper prepared for the International Conference on Better Health for Women and Children throughout Family Planning. Nairobi, Kenya; October 5-9.1987. New York: The Population Council.

[G] Bhatia, JC. Cleland, L Bhargavan and NSN Rao. 1997. Prevalence of Gynecological Morbidity. Among Women in South India. *Studies in Family Planning* 28(2):95-103.

[H] Deranisyagala, S. 1990. An Investigation into the Incidence and Causes of Domestic Violence Sri Lanka. *Women in Need (WIN)*, Colombo, Sri Lanka.

[I] Fortney, JA and BS Jason (eds) 1996. *The Base of the Iceberg: Prevalence and Perceptions of Maternal Morbidity in Four Developing Countries: The Maternal Morbidity Network*. Family Health International. Research Triangle, USA.

[J] Germain, A and R Dixon-Mueller, 1992. Stalking the Elusive 'Unmet Need for Family Planning. *Studies in Family Planning* 23(5):33-335.

[K] Gittlesohn, J, ME Bentley, PJ Pelto, S Nag, S Pachuri, A D Harrison and LT Landman, 1994. In: *Listening to Women Talk about their Health: Issues and Evidence form India*. New Delhi: Har-Anand Publications.

[L] Haq ul, M. 1991. *Human Development in South Asia*. Published for The Human Development Center. Karachi, Pakistan: Oxford University Press.

[M] Hawk, S. 1997. Why Include Men? Establishing STI clinics for Men in Rural Bangladesh. Health Policy and Planning (Submitted for publication).

[N] Indian Council of Medical Research. 1993, Uterine Cervical Dysplasia Study, Part II Program Report. New Delhi: Institute of Cytology and Prevention Oncology.

[O] International Institute of Population Studies. 1994. National Family Health Survey 1992-93, India Introductory, Report. Bombay.

[P] International Planned Parenthood Federation South Asia Region. 1994. Abortion in South Asia. IPPF-SAR. London (unpublished).

[Q] Jain, A and J Bruce. 1994. A reproductive health approach to the objectives and assessment of family planning programs. Pp.192-211 in Population Policies Reconsidered. Health Empowerment and Right, ed. G Lin. A Germain and L C Chen. Boston, USA. Harvard School of Public Health.

[R] Kapoor, I. 1997- Sexual and Reproductive Health Needs South Asia Region. Prepared for Consultation on Effective Mobilization and Use of Resources at International Parenthood Federation, South Asia Regional Bureau. New Delhi.

[S] Khan, ME, I Khan and N Mukherjee. 1997. Involving Men in Safe Motherhood. Journal of Family: Welfare 43(2):18-30.

[T] Koenig, M, S Jejeebhoy, S Singh and S Sridhar. 1996. Undertaking Community Based Research on the Prevalence of Gynecological Morbidity: Lessons from India. Prepared for IUSSP Seminar on Innovative Approaches to the Assessment of Reproductive Health Manila, The Philippines.

[U] Luthra, UK et al. 1992. Reproductive tract infections in India: the need for comprehensive reproductive health policy and programs. In Reproductive Tract Infections: Global Impact and Priorities for Women's Reproductive Health ed A Germain et al. New York and London: Plenum Press.

[V] Maine, D. 1991. Safe Motherhood Programs: Options and issues. Columbia University Center for Population and Family Health, New York.

[W] Menon, T, NS. Murthy, A Sehgal, AK Prabhakar and UK Luthra 1988. Role of Sexual Behavior in Cervical Cancer. Indian Journal of Preventive and Social

[X] Mishra, MK and TK Sinha. 1990. Cytological Screening for the Detection of Cancer in the Uterine Cervix. A Survey in Patna (India). *Cancer Lett* 52:21.

[Y] Murthy, MS, A Sehgal and L Satyanarayana. 1990. Risk Factors, Related to Biological Behavior of Precancerous Lesions of the Uterine Cervix. *British Journal of Cancer* 61:732-736.

[Z] National AIDS Control Organization. 1994. Country Scenario: An Update. Ministry of Health and Family Welfare, Government of India, New Delhi, India.

[AA] National AIDS Control Organization. 1997. NEXUS. Population Services International New Delhi, India.

[AB] National AIDS Control Organization. 1998. NEXUS. Population Services International. New Delhi, India.

[AC] Pachauri, S. 1994, Women's Reproductive, Health in India: Research Needs and Priorities. In *Listening to Women Talk about their Health Issues and Evidence from India*, ed. J Gittlesohn, ME Bentley, PJ Pelto, M Nag, S Pachauri, AD Harrison and LT Landman. New Delhi: Har-Anand Publications.

[AD] Pachauri, S. 1995. Defining a Reproductive Health Pack-age for India: A proposed Framework. South and East Asia Regional Working Paper No 4, New Delhi, India.

[AE] Pachauri, S. 1996. Reproductive Health: Concept, Ideology and Operational Issues. In *Perspectives in Reproductive Health* ed. J Sengupta and D Ghosh. New Delhi: New Age Publisher Ltd.

[AF] Pachauri, S. 1997. Breaking the silence. In *Whispers from Within: Unheard Voices of People with HIV/AIDS*, ed. S Solomon and RP Rajan. Navshakti Madras. India.

[AG] Population Action International 1995. Reproductive Risk : A World-Wide Assessment, of Women's Reproductive Health Data Sheet. Population Action International, Washington DC. USA.

[AH] Population Reference Bureau. 1996. The World's Youth Data Sheet. Population Reference Bureau, Washington DC. USA.

[AI] Rahman, F, M Whittaker and MR 1991. Maternal Mortality in Rural Bangladesh 1982-1990. Data from Verbal Autopsies. A presentation at ICDDR, B.

[AJ] Rodriguez. J, C Quintero. G Bergonzoli and A Salazar. 1985. Avoidable Mortality and Maternal in Cali Colombia. World Health Organization, Geneva (FHE/PMM/85.9.1) : 11.

[AK] Sarkar, S, S Panda , N Das, T Naik K Sarkar, BC Singh, JM Ralte, SM Ayer and SP Tripathi. 1993. Rapid Spread of HIV Among Injecting Drug Users in North-Eastern India. Indian Council of Medical Research, CARC CALLING 6:23-28.

[AL] Sidney, RS. SM Hashemi and AP Riley. 1997. Men's Violence Against Women in Rural Bangladesh: Undetermined or Exacerbated by Microcredit Programs? Paper presented at Population Association of America Annual Meetings Washington DC. USA.

[AM] Sinding, WS and M Fathalla. 1995. From Demographic Targets to Individual Needs: The Great Transition. Populi 22(8): 18-21.

[AN] Sood, S. (ed). 1990. Violence Against Women. Jaipur, India, Arihant Publishers.

[AO] United Nations. 1995. The World's Women 1995: Trends and Statistics. UN, New York, USA.

[AP] United Nations Children's Fund (UNICEF). 1996. The Progress of Nations 1996. New York.

[AQ] United Nations Development Program (UNDP). 1996. Report of the SAP Awareness Seminar. Quetta. 16 April 1996.

[AR] Walker. GJA, DEC Ashley, McCaw, and GW Bernard. 1986. Maternal Mortality in Jamaica. Lancet 1(8479) : 486-488.

[AS] Wasserheit, J et al. 1989. Reproductive Tract Infection in a Family Planning Population in Rural Bangladesh. *Studies in Family Planning* 29(2): 69-79.

[AT] World Bank. 1993. Investing in Health. World Development Report. New York: Oxford University Press.